


pSKAN Phagemid Display System

Product Information and Instructions
January 2001



Content

1. Background	4
2. Advantage of this system over other systems	7
3. Genetic information on the system	8
4. Protocols	11
4.1. Preparation and quantification of phage particles	12
4.2. Panning: selection of ligands	14
4.3. Characterization	16
4.5. Antibiotics, media and buffers	20
4.6. Bacterial strains	23
4.7. Antibodies	24
5. Literature	24
6. Examples, Applications	25
7. Order Information	31

The pSKAN Phagemid Display System was developed by Prof. Dr. J. Collins and Dr. P. Röttgen at the GBF Braunschweig, FRG. The system is patented.

A licence is required for the commercial use of any products developed with pSKAN.

The pSKAN Phagemid Display System

The pSKAN Phagemid Display System is an alternative to antibodies or immunoglobulin domain presenting phagemids. Since the pSKAN phagemid displays a small and constrained hypervariable amino acid loop, more than 3×10^7 variants can be tested for the specific binding to a target molecule of interest. The variants with the optimal (in general the highest) affinity can be selected and propagated without having to use animals or tissue cultures. The sequence of the binding domain can be determined easily by DNA sequencing. In comparison to antibodies, the pSKAN System is able to target very small epitopes, since it searches for binding sites with "only a fingertip" instead of "two whole hands" like antibodies. Within a period of only three weeks, the scientist is able to receive the target molecule of interest.

Applications:

- selection of peptides with desired binding affinities
- development of new affinity matrices for a protein of choice
- screening for novel protease inhibitors with potential clinical applications
- tools for the detection of proteins in e.g. food technology
- high throughput screening of protein-protein interactions
- selection of proteins with specific binding properties (high affinity binders)
- ligand screening
- development of pharmaceuticals and diagnostics
- detection of allergens
- of special interest: 3-D structure-based protein design
- and many more!

Advantages:

- can select very high affinity binders
- low dissociation due to the low flexibility and rigid structure of the hypervariable loop
- easy and reliable modelling because of the highly constrained epitope
- stability of the libraries due to the strong repression of the λ -promoter in *E. coli* WK6 λ mutS
- production of a soluble protein possible
- helpful for subsequent protein structure modelling



Mo Bi Tec

Advantages over immunoglobulins:

- ease of handling and rapidity (three weeks!)
- low cost
- animals or tissue cultures become superfluous

Advantages over immunoglobulin phagemids:

- targets smaller, recessed epitopes of limited accessibility
- monovalent display gives high selectivity

1. Background

The pSKAN Phagemid Display System can be used to select for and produce novel proteins which bind to target molecules of interest. Examples are the selection of new protease inhibitors (see chapter 6) or the creation of new peptide sequences which can be used like a monovalent antibody for an affinity purification.

The pSKAN Phagemid Display System produces molecular repertoires containing millions of site-specific mutant proteins in a form which enables an adsorption-based selection of both, gene and gene product. This property allows for repeated cycles of selection and amplification which results in the enrichment of novel proteins with the desired binding affinity for particular target molecules of interest.

The technology simultaneously selects variant genes and gene products. This is achieved by using the single-stranded bacteriophage M13 propagation system. The pSKAN phagemid containing a phage origin of replication is packed into a phage protein-coated particle on superinfection by a helper phage (M13K07 or derivatives). The phagemid encodes a hybrid protein consisting of a fusion between one of the phage coat proteins (pIII) and the human pancreatic secretory trypsin inhibitor (hPSTI) which presents a hypermutated amino acid loop (figures 1 and 2). This hPSTI-pIII_{M13} fusion protein replaces generally one (or two) of the five wild type pIII proteins of the phage coat, leading to a monovalent display of the hypervariable moiety (genetic details see chapter 3).

Originally, the hPSTI protein presents a loop of 7 amino acids stretched between two disulfide bridges on the protein surface. MoBiTec supplies three variations of this pSKAN library: due to the introduction of different oligonucleotides, the




Fig. 1. The infectious phagemid is displayed as an electron-microscopy picture (insert) and as a schematic drawing. The hybrid protein is shown enlarged as a drawing. Size of the phage: $60 \times 8800 \text{ \AA}$ ($0.006 \times 0.88 \text{ \mu m}$); size of the hPSTI protein: 6.1 kDa (production of a soluble hPSTI protein: see page 7).

pSKAN01/2001



Mo Bi Tec




Fig. 2:
The protein crystal structure of hPSTI in the van-der-Waals presentation: A: top view B: side view
pSKAN01/2001

loop consists of 6, 7 or 8 hypervariable amino acids. The three libraries are referred to as HyC, HyB and HyA, respectively (Hy for hypervariable; HyC has a 7 amino acid loop of which only 6 are hypervariable). By randomly mutagenizing the hPSTI gene sequence for these amino acid loops, three different libraries of phagemids are constructed containing tens of millions of ligand variants. Some of these gene product variants will have the desired binding affinities for the particular target molecule of interest. The subset of phagemids with the desired binding properties is selected by, for example, binding on a surface coated with the target molecule; the poorly binding majority of phagemids being washed away. The strongly binding population is eluted and amplified by reinfection and cultivation in *E. coli*. About three successive rounds of such binding and amplification lead to the enrichment of one (or very few) variant ligands which exhibit the desired binding characteristics for the target molecule. The three libraries are normally treated separately during the selection process, but a combination of the libraries is possible as well.

After selecting the peptide sequence of interest, the production of a soluble hPSTI protein is possible as well. For this purpose, the DNA sequence of the selected hypervariable loop can be cloned into a vector like pMAMPF (reference #4). pMAMPF vectors produce a soluble variant of the hPSTI protein. By exchanging the DNA sequence of the hPSTI loop in pMAMPF with the DNA sequence of the selected pSKAN phagemid, a small soluble protein of 6.1 kDa can be produced in *E. coli*.

2. Advantage of this system over other systems (Immunoglobulins, etc.)

The pSKAN Phagemid Display System represents an alternative to the peptide or peptide analogue libraries and the RNA evolutive technology in the search for novel ligands. Different from these latter systems, the pSKAN phagemid presents the variable epitope under defined conformational restraints on the protein surface between two disulfide bridges. Over a linear cross-section of 30-40 Å a binding site of 5-7 amino acids is displayed without problems of steric hindrance from neighbouring molecules, allowing targetting of specific regions. The crystal and solution (NMR data) structures of hPSTI and a number of variants are known, so that the structure of new variants, which differ only in the loop, can be accurately estimated and modelled. The selected variants can serve as excellent "lead molecules" for the development of synthetic mimotopes.



Advantage over other phage display systems:

As mentioned in chapter 1., the pSKAN phagemid presents only one or two hybrid hPSTI-pIII_{M13} molecules. Such essentially monovalent phagemids exhibit a strong selective advantage for variant molecules with high avidity. In phage display systems, where the pIII gene is replaced on the DNA level by a recombinant gene producing the hybrid protein, the fusion protein replaces all five pIII molecules. In such a system, phages which display molecules with less affinity will also bind (via five variable moieties) to the target molecule of interest resulting in the selection of a less specific variant molecule.

Advantage over immunoglobulins:

Compared to immunoglobulins, the pSKAN phagemid has the following advantages: The development and production of polyclonal and monoclonal antibodies is time-consuming. Furthermore, animals (mostly mice) have to be injected and finally are normally killed. For polyclonal antibodies this process has to be repeated for every new production. For monoclonal antibodies, the selected cell lines have to be kept, grown and harvested for production, which is a rather expensive process. In addition, antibodies are large molecules consisting of four peptide chains. Thus, once parts of the molecules unfold, refolding is rather improbable. The large antibodies try to find a marked epitope of a molecule with two binding sites ("with two hands") while the pSKAN phagemid searches for an epitope with a small amino acid loop ("with a fingertip"). Antibodies have problems targetting small epitopes due to steric hindrance by their surface peptide loops. The hPSTI protein on the contrary is small and consists of only one peptide chain. Compared to antibodies, the pSKAN Phagemid Display System is inexpensive, easy to handle, does not require involved laboratory equipment and surveillance and results in a variant protein within a very short period of time.

Advantage over immunoglobulin phagemids:

The pSKAN Phagemid Display System also has advantages over immunoglobulin domain-presenting phagemids for "screening" small epitopes. In comparison to F_{ab} phagemids (hypervariable CDR3 region), the varied epitope of pSKAN is small and constrained as an extended peptide. The immunoglobulin phagemids only display up to 10⁵ variations, a small number compared to the 3x10⁷ variants displayed by the pSKAN system.

Clinical potential:

hPSTI is based on a small human acute response protein (human pancreatic secretory trypsin inhibitor) which is present in high levels in serum. This might be of advantage at a later stage, if direct clinical application is envisaged for the variant protein (e.g. development of new protease inhibitors with therapeutic potential; see chapter 6.2.).

3. Genetic information on the system

The pSKAN phagemid, which combines genetic elements from phage and plasmid, uses the ColE1 plasmid origin for replication in *E. coli*. Upon superinfection with the helper phage M13K07, single stranded DNA is produced from the phage fd origin. The phagemid DNA is then packaged into the M13K07 phage coat, creating a new phagemid generation with pSKAN-DNA. A hybrid protein consisting of a fusion between one of the M13 phage coat proteins (pIII) and the human pancreatic secretory trypsin inhibitor (hPSTI) is expressed from the phagemid DNA. This fusion protein is regulated by the λP_L promoter. Thus, it is repressed in the λ -lysogen *E. coli* WK6 λ mutS which produces the λ -repressor cl. hPSTI-pIII_{M13} is only expressed after superinfection with the helper phage. The induction mechanism, which is initiated by the superinfection, is not fully understood.

hPSTI-pIII_{M13} can replace one or more of the five wild type pIII_{M13} coat proteins during phage coat assembly. Statistically, only one hybrid protein is integrated in every second phagemid. Thus, a population of phages is produced, which monovalently displays the hybrid protein with the hypervariable loop (see figures 1 and 2) and which contains pSKAN phagemid DNA. When the new phagemid generation infects *E. coli* again, the phagemid DNA remains in form of a plasmid in the cell and no further phages are produced. For phagemid production, a new infection with helper phage is essential (see figure 3).

Since the helper phage is a M13 derivative, the cells are not lysed; the phages are excreted. The host strain needs to carry the F-episome encoding the sex pili which serve as phage receptors during the infection.




Fig. 3: pSKAN Phagemid Life Cycle in *E. coli* (squares)





Fig. 4: Phagemid vector construction. The phagemid contains the genetic elements: selection marker Amp^R (ampicillin resistance); $ColE1$ ori: plasmid origin of replication; fd ori: origin of replication for the fd phage; λp_L : λ -promoter left; $ompA$: signal sequence for secretion; $hPSTI$: gene of the human pancreatic secretory trypsin inhibitor; the $hPSTI$ gene contains the inserted randomized sequence $(NNG/T)_n$ with n being 6 (HyC), 7 (HyB) or 8 (HyA), Hy for hypervariable; $pIII$: gene for the phage coat protein III; $2 \times t_{fd}$: two phage fd transcription terminator sequences; cat_{amber} : chloramphenicol resistance is inactive due to an amber mutation.

The complete phagemid sequence of the original vector pSKAN8, which was used for the creation of the libraries, is available for download from our web sites <http://www.mobitec.de> or www.mobitec-us.com and is accessible in the EMBL data base, accession # X84307. pSKAN8 is based on the vector pMAMPF-3 (reference 1).

pSKAN01/2001

4. Protocols

Note: MoBiTec conveniently provides the pSKAN libraries HyA, HyB and HyC ready-to-use in three 1 ml aliquots of *E. coli* WK6λmutS. The host strain (without phagemid) for re-infection is supplied with the libraries. Therefore, after the preparation of the phagemids, the customer is immediately able to start the selection of ligands, the so called "panning" procedure. If there is an interest in preparing your own custom-made pSKAN DNA library. The original pSKAN8 vector, which was also used to prepare the above mentioned libraries, is available at MoBiTec as well (see chapter 7).



*HyA (n=8), HyB (n=7), HyC (n=6); n indicating the number of hypervariable amino acids in the loop



Mo Bi Tec

All buffer compositions, media and antibiotics are listed in chapter 4.5.

An example suitable as positive control is described in chapter 6.1.

We highly recommend the performance of this positive control!

General knowledge: Users of this kit should be familiar with molecular biology and microbiological techniques!

4.1. Preparation and quantification of phagemid particles

Note: Since every amplification of a library causes the loss of peptide variants, we strongly recommend to work with the original library stocks provided by MoBiTec which has 10^7 variants per library. The host strain *WK6 λ mutS* without phagemid (for re-infection) should be kept on minimal medium during cell keeping in order to select for the F-episome, which carries proAB for the complementation of the Prolin synthesis deficiency.

4.1.1. Helper phage: M13K07 phage stocks

The preparation of M13K07 helper phages should be started from a single fresh phage plaque. Therefore, inoculate 20 ml of LB medium with a single colony of *E. coli* WK6 cells and incubate over night at 180 rpm and 37°C. Use 200 μ l of this culture to inoculate 20 ml LB medium and incubate at the same conditions until the culture reaches the logarithmic growth phase (2-3 hours; $OD_{600}=0.5$). Mix 1 μ l of a M13K07 phage stock solution (e.g. from Pharmacia) and 0.5 ml of logarithmic growing WK6 cells with 3 ml of molten LB top agar (about 40°C) and pour the mixture onto a LB agar plate. Incubate over night at 37°C. The next day, use a disposable Pasteur pipette to pick a single, well separated phage plaque and inoculate 20 ml of LB (2x)/Km medium (100 ml Erlenmeyer flask). Incubate overday at 37°C on a shaker at 180 rpm. Inoculate 2 x 500 ml LB (2x)/Km medium with 10 ml preculture and incubate overnight (37°C, 180 rpm).

The next day, centrifuge four 250 ml aliquots for 15 minutes at 8,000 rpm and 4°C (GS3 rotor, Sorvall RC5C). Transfer the supernatant into centrifuge bottles and centrifuge again. Transfer the supernatant again, add 0.15 vol of PEG/NaCl solution, mix and incubate on ice for at least 2 hours.

pSKAN01/2001

Centrifuge for 40 minutes at 8,000 rpm (GS3 rotor), decant the supernatant, repeat the centrifugation for 1 minute at 4000 rpm and remove last traces of supernatant using a pipet. Resuspend each PEG-pellet in 2.5 ml PBS solution and collect the resuspended phages in one SS34 centrifuge bottle. To clear the suspension centrifuge again for 10 minutes at 12,000 rpm (SS34 rotor). Recover the supernatant (pipet), add NaN_3 to a final concentration of 0.02% and store the phages at 4°C.

Note: About three to four successive rounds of binding and amplification lead to the enrichment of one (or very few) variant ligands which exhibit the desired binding characteristics for the target molecule. Therefore, the following protocols have to be repeated at least three times.

4.1.2. Packaging of pSKAN phagemids

Inoculate 20 ml LB medium containing 250 µg/ml Amp and 20 µg/ml Tc in a 200 ml Erlenmeyer flask with 200 µl from the frozen *E. coli* WK6λ.mutS stocks containing the pSKAN phagemids (as provided in the MoBiTec kit or from the previous panning step). Incubate overnight at 37°C.

Note: An inoculation with 200 µl provides more than sufficient cells than possible peptide variants (10^7 per library). The concentration (cells/ml) for each lot is indicated on the data sheet provided with the product.

For packaging the phagemids, inoculate 100 ml of LB/Amp/Tc medium (1 l Erlenmeyer flask) with 1 ml of the overnight culture. Incubate at 37°C and 180 rpm until $\text{OD}_{600} = 0.5$ (~ 2.5 hours). Add 500 µl of the helper phage M13K07 stock solution (10^{11} - 10^{12} cfu/ml; see 4.1.1.), incubate at 37°C without shaking for 15 minutes and then at 37°C and 180 rpm over night. *Note:* Some scientists have observed higher phagemid yields, when Kanamycin (end concentration 50 µg/ml) was added 0.5 to 1 hour after infection with the helper phage, leaving sufficient time for the expression of the helper phage's Km resistance gene.

The next day, centrifuge 10 minutes at 8,000 rpm (GSA rotor), decant the supernatant into a fresh bottle and repeat the centrifugation step. Add 0.15 vol of PEG/NaCl solution to the supernatant and incubate on ice for at least two hours. Centrifuge for 30 minutes at 10,000 rpm (GSA rotor), decant the supernatant and repeat the centrifugation. Remove the supernatant completely, dissolve the pellet in 1 ml of PBS buffer and transfer the solution into an Eppendorf reaction tube. Centrifuge for 10 minutes at 13,000 rpm (batch centrifuge) recover the cleared solution and add NaN_3 (final concentration of 0.02%). Store at 4°C.



Mo Bi Tec

4.1.3. Determination of phagemid titers (colony forming units [CFU] assay)

Inoculate 20 ml of LB/Tc (20 µg/ml) medium with 200 µl of an *E. coli* WK6λmutS overnight culture (37°C; LB/Tc) and incubate at 37°C and 180 rpm for 2 to 3 hours (OD₆₀₀ = 0.5).

For each phagemid probe fill seven to eight wells of a sterile 96-well culture dish with 90 µl of autoclaved water and prepare dilution series by transferring 10 µl aliquots (dilutions 10⁻¹ to 10⁻⁷ or 10⁻³ to 10⁻¹⁰).

Add 100 µl of logarithmic growing cells, mix and incubate for 30 min at 37°C. Spot 20 µl portions of each well on LB/Amp/Tc (phagemid titer) and LB/Km/Tc (helper phage titer) agar plates and incubate over night at 37°C. As a control use 20 µl of noninfected log-phase cells spotted on LB/Amp/Tc, LB/Km/Tc and LB/Tc agar plates.

4.2. Panning: selection of ligands

Note: Several panning procedures can be applied at this stage. The protocols below describe the standard procedure with microtiter plates for the coating step which is used at the GBF Braunschweig, FRG. Alternative coating protocols involve tubes, magnetic beads or the concentration of the ligands. Variations are also possible - and advisable - at the elution step. An example for phagemid titers resulting from a panning procedure is given in chapter 6.1. It is recommended to perform this example as positive control.

4.2.1. Panning procedure

4.2.1.1. Coating of microtiter plates

Fill 100 µl of ligand solution (100 µg/ml PBS) into the wells of a 96-well microtiter plate (Nunc, Maxisorb; has a hydrophilic surface for the binding of ligands). Incubate over night at 4°C or at least 2 hours at room temperature.

Note: One well is sufficient per ligand and per reaction condition (e.g. different elution buffers; see figure 6). One well should be left without ligand as a negative control. If phagemids are enriched in this negative control after 3-5 rounds, they might have an affinity to the blocking reagent (skim milk). In such a case, it is advisable to start a new panning procedure.




Fig. 5: General application protocol

pSKAN01/2001

Mo Bi Tec

Shake out the wells, slap the plate onto a paper towel and wash the wells once with T-PBS solution (ELISA plate washer).

4.2.1.2. Blocking

In order to block any free positions, fill the wells with 400 μ l of blocking solution and incubate at room temperature for \sim 1 hour. Shake out the wells, slap the plate onto a paper towel and wash the wells once with T-PBS.

4.2.1.3. Binding

Fill the coated and one uncoated well (as a negative control) with 100 μ l of phagemid preparations diluted 1:1 with blocking solution or PBS ($\sim 10^{10}$ - 10^{11} phages/well) and incubate at room temperature for at least 1 hour.


4.2.1.4. Washing

Shake out the wells and slap the plate onto a paper towel. In the first round of panning wash the wells once with T-PBS, incubate for 10 minutes with 400 μ l blocking solution, wash again with T-PBS and finally two times with water. During all further rounds repeat the T-PBS washing steps three times. All washing steps may be carried out by using an ELISA plate washer.

4.2.1.5. Elution

Slap out the plate and fill the wells with 100 μ l of elution-buffer, incubate at room temperature for 15 minutes and transfer the solution into an Eppendorf reaction tube containing 6 μ l Tris (2 M).

Determine the titer of eluted phages as described under 4.1.3.



*1-4 indicating different reaction conditions (e.g. variation in the elution buffer)

Fig. 6: Panning with microtiter plates. Per reaction condition (#1-4) one well per library (HyA, HyB, HyC) is coated with the ligand of interest. After 3-5 rounds an increase in the pSKAN phage titer is observed. The control without panning target shows the background of unspecifically binding phages. The helper phage titer (M13K07) serves as additional control. Km, kanamycin; Amp, ampicillin.

pSKAN01/2001

4.2.2. Reinfection of *E. coli* cells

Mix the eluted phages and 10 ml of *E. coli* WK6 λ mutS log-phase cells (37°C; LB/Tc- culture) and incubate for 30 min at 37°C. Collect the cells by centrifugation (5 minutes, 8,000 rpm, SS34 rotor) and resuspend the pellet in 400 μ l of LB/Amp (250 μ g/ml)/Tc (20 μ g/ml) medium. Plate 200 μ l aliquots onto LB/Amp/Tc agar plates and incubate them overnight at 37°C.

4.2.3. Packaging of phagemids from reinfected cells

Resuspend the complete lawn of the reinfected *E. coli* cells in 20 ml of LB/Amp/Tc medium and use 2 ml for inoculation of 50 ml LB/Amp/Tc medium (250 ml Erlenmeyer flask). Incubate at 180 rpm and 37°C for 1 hour, add 100 μ l of M13K07 stock solution (10^{11} - 10^{12} cfu/ml; see 4.1.1.) and incubate for 15 minutes at 37°C without shaking. Continue the incubation at 37°C and 180 rpm overnight.

The next day, collect the phages as described under 4.1.2.

After each round of panning, the titers of kanamycin resistant (KmR; helper phage) and ampicillin resistant (AmpR; pSKAN) phagemid particles were determined according to 4.1.3.

Note: An increase of eluted pSKAN phages should be recognized after 3-5 rounds (see figure 6). Example titer results are given in chapter 6.1.2.

4.3. Characterization

4.3.1. DNA Sequencing

Note: After three to four rounds of panning, a first characterization of the selected phagemids is advisable. For this purpose, sequencing the single clones is the easiest procedure.

Use standard protocols for DNA sequencing. Suitable primers are available at MoBiTec (see chapter 7). The primer sequences are as follows:

- for single- and double-stranded DNA (binding site at the start of pIII; sequencing towards hPSTI):

primer # 1255 5' GGGATTTTGCTAAACAAC 3'

- for double-stranded DNA (esp. Alf-sequencing; binding site before hPSTI leader sequence; sequencing towards hPSTI):

primer # 2897 5' GGAGGTCTAGATAACGAGG 3'


Note: We recommend double-stranded sequencing, since primer # 1255 also recognizes a sequence on the helper phage M13K07, which would be added for the production of single-stranded DNA.

4.3.2. Protein Gels and Western Blots

Note: Protein gels and Western blots are used as a control for the integration of hybrid proteins in the phagemid particles. An anti-pIII antibody is available at MoBiTec (see chapter 4.7. and 7). Alternatively, an hPSTI antibody is available (see chapter 4.7.) which can be used to detect hPSTI directly. The antibody is also suitable to detect soluble hPSTI. In a Western blot the hPSTI protein yields one band at 6.1 kDa. Phagemids containing a hPSTI-pIII hybrid protein will yield two bands in a Western blot, one resulting from the non-hybrid pIII proteins, the other from the larger fusion protein hPSTI-pIII. Phagemids without hybrid protein yield only one band (for pIII). Such phagemids can occur, when a stop codon is present within the selected hypervariable loop. Cells without a fusion protein grow faster and could thus potentially be enriched. This control will exclude a false positive result.

Use standard protocols for polyacrylamide gel electrophoresis (PAGE) of proteins, blotting and protein detection by specific antibodies.

Fig. 7: Schematic presentation of a Western blot using the pIII antibody. A phagemid displaying a hybrid protein (such as pSKAN8 or a phagemid selected during the panning procedure) shows two bands, one for pIII and one for the larger hybrid protein hPSTI-pIII. A phage, which does not present a hybrid protein, like M13K07 or a pSKAN phagemid with a stop codon within the hypervariable region, shows only one band.



pSKAN01/2001

4.3.3. Phagemid ELISA

Note: In order to test the binding capacity of the isolated phagemid variants an ELISA can be used. An example of an ELISA result is given in chapter 6.1. (figure 8).

To coat a microtiter plate, fill 100 μ l solution of the desired ligand (e.g. 100 μ g/ml PBS) into the wells and incubate at 4°C overnight.

Slap out the wells and wash them once with T-PBS solution. Add 400 μ l of blocking solution and incubate for 1 hour at room temperature and wash again. Incubate with phages (4.2.2.) diluted 1:1 in blocking solution for 3 hours and wash two times with T-PBS. Incubate with a commercially available anti M13 antibody (anti pVIII, peroxidase labeled; diluted 1:5,000 in blocking solution; available from Amersham Pharmacia Biotech) for 1 hour and wash the wells three times with T-PBS. Incubate with 100 μ l of staining solution, stop the reaction after 30 minutes by adding 50 μ l of 2 M H₂SO₄ and measure the absorption at 492 nm.

To exclude any false positiv results, the following negative controls are recommended for the phagemid ELISA:

- a) well without ligand (to exclude unspecific phagemid binding to the plate or the blocking reagent)
- b) well without phages (to exclude, that the M13 antibody binds unspecifically)
- c) well without M13 antibody (as a control for the antibody-specific color reaction)

4.5. Antibiotics, media and buffers

Antibiotics (stock solutions)

Ampicillin (Amp 50 mg/ml):

Dissolve 1g of ampicillin (sodium salt) in 20 ml water and sterilize by filtration

Tetracycline (Tc, 20 mg/ml):

Dissolve 400 mg of tetracycline in 20 ml of ethanol (50%)

Kanamycin (50 mg/ml):

Dissolve 1 g of kanamycin in 20 ml of H₂O
and sterilize by filtration.

Store all antibiotics at -20°C

M9 medium (1x)

M9 salt solution (10x)	100 ml
glucose (40% w/v)	12.5 ml
CaCl ₂ (1 M)	100 µl
MgSO ₄ (1 M)	1 ml
FeCl ₃ (1 mM)	0.5 ml
Thiamine (10 mg/ml)	100 µl
H ₂ O	ad 1 l

all solutions are autoclaved or sterile filtered (Thiamine) separately

M9 salt solution (10x)

Na ₂ HPO ₄ *2 H ₂ O	74.1 g
KH ₂ PO ₄	30 g
NaCl	5 g
NH ₄ Cl	10 g
H ₂ O	ad 1 l
autoclaved	

LB stock solution (2x)

10 g tryptone
10 g yeast extract
5 g NaCl
H₂O ad 500 ml
autoclaved and stored at 4°C

Mo Bi Tec

Agar stock solution (2x)

15 g agar
H₂O ad 500 ml
autoclaved and stored at 4°C

LB medium

10 g tryptone
10 g yeast extract
5 g NaCl
H₂O ad 1,000 ml
autoclaved

LB agar plates

For preparation of 40 agar plates melt 500 ml of agar stock solution, add 500 ml of LB stock solution, mix and pour into sterile petri dishes.

For selective plates add 5 ml of ampicillin (50 mg/ml), 1 ml of tetracycline (20 mg/ml) or 1 ml of kanamycin (50 mg/ml) stock solution respectively.

LB top agar

LB medium containing 0.6% Agar

PEG/NaCl-solution: (16.7%/3.3 M)

100 g PEG 8,000
116.9 g NaCl
475 ml H₂O

PBS-buffer (1x, pH 6.8 - 7.0):

8.0 g NaCl
0.2 g KCl
1.43 g Na₂HPO₄ + 2 H₂O
0.2 g KH₂PO₄
H₂O ad 1 l

For panning and phagemid ELISA:

T-PBS solution:

PBS-buffer containing 0.5% Tween 20

Blocking solution:

PBS-buffer containing 2% skim milk powder

Elution-buffer:

glycine (0.1 M; pH 2.2)

staining solution for phagemid ELISA:

3.1 ml Na_2HPO_4 (0.2 M)

2.9 ml citric acid (0.2 M)

6 ml H_2O

4.8 mg o-phenylenediamine

1.5 μl H_2O_2 (30%)

4.6. Bacterial strains

- *E. coli* *WK6λmutS* cells with and without the pSKAN libraries (supplied by MoBiTec)

Genotype *E. coli* *WK6λmutS*: *galE*, *strA*, *nalT*, (*lac-proAB*), λ^+ , *mutS::Tn10*, *F'* [*lac^q*, *lacZ* M15, *proAB*⁺];

The host strain *WK6λmutS* without phagemid (for re-infection) should be kept on minimal medium (M9) during cell keeping in order to select for the F-episome, which carries *proAB* for the complementation of the prolin synthesis deficiency. The transposon Tn10 carries the Tc resistance gene. The original strain *WK6mutS*, which does not contain the λ lysogen, is described in reference 13.

- for the helper phage: *E. coli* *WK6* or any *E. coli* *F*⁺-strain (not provided) (M13 and derivatives require the presence of an F-episome, since the here encoded sex pilus serves as phage receptor.)

4.7. Antibodies

- anti-pIII (g3p) antibody
mouse monoclonal IgG, unlabeled; use 1:1000 for Western blots
(see chapter 4.3.2. and reference 14).
- anti-hPSTI antibody
mouse monoclonal IgG, unlabeled, use 1:1000 for Western blots and ELISA.
Recognizes soluble hPSTI as well as the hPSTI in the pIII-hPSTI fusion protein
of pSKAN.

5. Literature

1. Maywald, F. *et al.*, *Gene* 68, 357-369 (1988)
2. Böldicke, T. *et al.*, *Eur. J. Biochem.* 175, 259-264 (1988)
3. Collins, J. *et al.*, *Biol. Chem. Hoppe Seyler* 371, 29-36 (1990)
4. Szardenings, M. and Collins, J., *Gene* 94, 1-7 (1990)
5. Paulsen, J. *et al.*, *Ber. Bunsenges. Phys. Chem.* 93, 1015-1019(1989)
6. Paulsen, J. *et al.*, *DECHEMA Biotech. Conf., DECHEMA, Frankfurt*, pp1031 (1989)
7. Hecht, H. J. *et al.*, *J. Mol. Biol.* 220, 711-722 (1991)
8. Collins, J. *et al.*, in *Wooley et al. (ed.) D.F.I. Print, Aarhus*, pp 34 (1990)
9. Klaus, W. and Schomburg, D., *J. Mol. Biol.* 229, 695-706 (1993)
10. Hecht, H. J. *et al.*, *J. Mol. Biol.* 225, 1095-1103 (1992)
11. Szardenings, M. *et al.*, *Protein Engineering* 8, 45-52 (1995)
12. Röttgen, P. and Collins, J. , *Gene* 164, 243-250 (1995)
13. Zell, R. and Fritz, H.-J., *EMBO J.* 6, 1809-1815 (1987)
14. Tesar, M. *et al.*, *Immunotechnology* 1, 53-64 (1995)
15. Dimasi, N. *et al.*, *J. of Virology*, 71, (No.10) 7461-7469 (1997)
16. Mersmann, M. *et al.*, *J. Immunol. Meth.* 220, 51-58 (1998)

6. Examples, Application

6.1. Positive control: Selection and characterization of chymotrypsin binding hPSTI-variants

The following application example describes the use of one of the supplied phagemid libraries for the selection of protein variants with altered binding specificity, namely novel chymotrypsin inhibitors (see also reference 12). **We highly recommend to perform these experiments as positive controls.**

The phagemid pSKAN8, which was used for the construction of the pSKAN phagemid libraries HyA, HyB and HyC, encodes a hPSTI-pIII fusion protein which is a specific inhibitor of the protease trypsin. pSKAN8 can be purchased separately (see chapter 7).

6.1.1. Methods

Preparation of phagemid particles

Phagemids were packaged from library HyB according to 4.1.2. using 3×10^{11} (Km^{R} cfu/ml) M13K07 helper phages. The final preparation contained 2×10^{12} Amp^{R} cfu/ml and 8×10^{10} Km^{R} cfu/ml (determined according to 4.1.3.).

Panning procedure

Coating (according to 4.2.1.1.)

One well of a 96-well microtiter plate (Nunc, Maxisorb) was filled with 100 μl of α -chymotrypsin (bovine; 100 $\mu\text{g}/\text{ml}$ PBS) and incubated over night at 4°C. The next day the plate was washed twice with T-PBS.

Blocking

The chymotrypsin coated well and an additional uncoated well (to be used as a negative control) was blocked according to 4.2.1.2.

Mo Bi Tec

Binding (according to 4.2.1.3.)

100 µl of phagemid preparations were pipetted in the coated and uncoated wells (both blocked) and incubated for 3 hours on a rocker at room temperature.

For the first cycle the phagemid preparation mentioned above was diluted by a factor of 10 in blocking solution. For all of the successive cycles phages prepared from reinfected cells (see below) were diluted by a factor of 2. For exact titers see table 2.

Washing

The washing procedure (according to 4.2.1.4.) is summarized in table 1. All washing steps were performed using a Titertek Mikroplate Washer S8/12 (Flow Laboratories GmbH, Meckenheim, FRG), superwash mode, 600 µl washing solution per well.

Table 1: Washing procedure

procedure	round 1	round 2	round 3 to 5
washing with T-PBS	2x	3x	5x
incubation with blocking solution	10 min.	10 min.	10 min.
washing with T-PBS	2x	3x	5x
incubation with blocking solution	5 min.	5 min.	5 min.
washing with T-PBS	2x	3x	5x
washing with H ₂ O	2x	2x	2x

Note: this washing procedure has been proven as very selective. In some cases a less stringent procedure may be necessary.

Elution: according to 4.2.1.5.

Re-infection and packaging of phagemids from re-infected cells.

The eluted phages were used for re-infection of *E. coli* cells as described under 4.2.2. and plated separately.

Resulting bacterial lawns were resuspended and phagemid particles for the next round of panning were prepared according to 4.2.3.

6.1.2. Results

Enrichment of chymotrypsin binding variants

The panning experiment was done for a total number of five cycles. After each round of panning the titers of kanamycin resistant (Km^R ; helper phage) and ampicillin resistant (Amp^R ; phagemid containing) phage particles were determined according to 4.1.3. for both input solutions and eluates (table 2).

Table 2: An example of phage titers during an anti-chymotrypsin panning

panning target		round 1	round 2	round 3	round 4	round 5
input						
α -chymotrypsin						
Amp^R	2×10^{11}	1×10^{11}	1×10^{11}	3×10^{11}	1.5×10^{11}	
Km^R	8×10^9	3×10^{10}	7.5×10^9	4×10^9	n.d.	
none						
Amp^R	2×10^{11}	4×10^{11}	3×10^{11}	5×10^{11}	1×10^{10}	
Km^R	8×10^9	8×10^9	1×10^{10}	1×10^{10}	n.d.	
eluate						
α -chymotrypsin						
Amp^R	2×10^4	5×10^4	4.5×10^6	2×10^7	2×10^7	
Km^R	n.d.	n.d.	5.4×10^4	1×10^5	2×10^6	
none						
Amp^R	2×10^5	5×10^4	4.3×10^4	7×10^4	4×10^4	
Km^R	n.d.	n.d.	1.5×10^3	3×10^3	2×10^4	

Titers are given as ampicillin resistant (Amp^R) or kanamycin resistant (Km^R) colony forming units (cfu) per ml; n.d.: titer not determined.

Mo Bi Tec

Explanation

As can be seen from table 2 the titer of input phages were roughly the same in all rounds of panning in both cases. In the case of chymotrypsin an increase of eluted phages can be definitively recognized for the first time after the third round (4.5×10^6 vs. 4.3×10^4 Amp^R cfu/ml). This leads to a further, stable enrichment during round four and five (2×10^7 Amp^R cfu/ml). The background of unspecific binding phages (ca. 5×10^4) is represented by the negative control (without target).

Characterization of enriched variants

DNA-Sequencing:

The enrichment of chymotrypsin-binding variants within the eluted phage population was verified by DNA-sequencing of the hypervariable region of randomly selected single clones. (Normally ten single clones were taken from the titration plates after round three to five and sequenced using primer # 2897.)

Regarding the chymotrypsin-panning, only three different clones (Chy1, Chy4 and Chy5*) were found among six random clones from round three. An additional clone (Chy8*) appeared in round four. The predominant clone of round three was further enriched to 80% (8/10) and 100% (10/10) of the clones sequenced after round four and five. All of these clones show homology to each other as well as to known chymotrypsin inhibitors.

Within the control an enrichment of single clones could not be detected during rounds three to four. All of the clones sequenced were clearly unrelated in sequence and 30% to 60% of them contained stop codons.


*for sequences see Röttgen and Collins, (reference12, page 24)

Phagemid ELISA

To verify that the isolated variants are able to bind their target, phagemid particles were used for an phagemid ELISA according to 4.3.3. For results, see figure 8, page 29.

Figure 8 shows, that all of the isolated clones are indeed chymotrypsin binders and show no affinity for closely related targets like trypsin or unspecific binding to the coated surface.

Fig. 8: ELISA of chymotrypsin binding phages
 Each row was coated with the target proteins noted on the right and each column was incubated with phage particles noted below. M13K07: helperphage particles; pSKAN8: phages presenting the original hPSTI-variant; Chy-1 to Chy-8: phage clones isolated by anti-chymotrypsin panning. The phage titers of the original preparations are noted in brackets and dilutions of these used per row are given as phage dilution factors (pdf).



pSKAN01/2001

Concluding remark

The experiment described here was one of a series of panning experiments done at the GBF, which demonstrated the usefulness of the pSKAN phagemid display system for affinity selection approaches.

The data presented here may serve as a clue for your own panning experiments. Corresponding experiments may also be included as positive controls.

6.2. Highly effective protease inhibitors from variants of human pancreatic secretory trypsin inhibitor (hPSTI): an assessment of 3-D structure-based protein design (reference 11, page 24)

Variants of hPSTI were generated in order to design extremely strong and specific human leukocyte elastase (HLE) inhibitors. HLE is involved in several disorders, like adult respiratory distress syndrome (ARDS) or septic shock. Thus, a HLE inhibitor could have potential therapeutic applications. Szardenings *et al.* (reference 11) were able to develop an excellent inhibitor with a minimum of two amino acid exchanges.

Protein purification:

0.05% Tween-80 and 5.5% perchloric acid were added subsequently to the supernatant of a typical 1 l batch fermentation. The precipitate was removed by centrifugation after 30 minutes and the solution re-adjusted to pH 8.0 by the addition of a saturated solution of Tris. Salts were removed by dialysis against water in a dialysis tubing (3 kDa cut-off). Further purification was carried out by filtration through a DEAE Sephacel ion exchanger. The flow-through and the first fractions obtained by washing with 20 mM Tris pH 8.0 were adjusted to pH 2.7 with citric acid and applied to an S-Sepharose fast flow column. The column was washed with 10 mM citric acid (pH 2.7) and most hPSTI variants could be obtained from a NaCl gradient (0 - 1 M NaCl) at about 0.6 M NaCl. Depending on batch and particular variant, the hPSTI already represented about 50% of the entire protein content of the peak fractions (judged by SDS PAGE). The protein was either purified further by an additional chromatography on a Mono-S column or the fractions were dialysed against water and lyophilized. The final purification was carried out by reverse-phase chromatography on a ProRP column running a gradient from 0 to 60% acetonitrile containing 0.1% TFA.

pSKAN01/2001

Note: For such an application, the production of soluble protein is essential, i.e. the protein has to be produced without the pIII-portion. Therefore, the DNA sequence of interest has to be cloned into another vector before protein production. (Please see pMAMPF vectors described in reference 4)

7. Order Information

order #	description	amount
store at -70°C:		
PSKAN	3 pSKAN libraries in <i>E. coli</i> incl. <i>E. coli</i> host strain <i>WK6λmutS</i>	4 x 1ml
store at +4°C:		
PS1255	sequencing primer # 1255 (18 mer) lyophilized	500 pmole
PS2897	sequencing primer # 2897 (19 mer) lyophilized	500 pmole
pSKAN8	pSKAN8, lyophilized vector DNA	5μg
PSKAN4	pMAMPF-3-hPSTI4, lyophilized vector DNA	5μg
store at -20°C:		
PSKAN2	anti-hPSTI antibody, monoclonal mouse lyophilized	40 μg
PSKAN3	anti-pIII (g3p) antibody, monoclonal mouse lyophilized	100 μl